arXiv:quant-ph/9903004v2 7 Apr 1999

Entanglement degree for Jaynes-Cummings model

S. Furuichi
Science University of Tokyo in Yamaguchi,
1-1-1 Daigakudori Onoda city Yamaguchi, 756-0884, Japan

M. Ohya
Department of Information Sciences, Science University of Tokyo,
2641 Yamazaki Noda city Chiba, 278-8510, Japan

(February 1, 2008)

Recently, it has been known that a quantum entangled state plays an important role in the field of quantum information theory such as quantum teleportation and quantum computation. The research on quantifying entangled states has been done by several measures. In this letter, we will adopt the method using quantum mutual entropy to measure the degree of entanglement on the Jaynes-Cummings model.

PACS numbers: 89.70.+c, 03.65.Bz, 42.50.Dv

The degree of entanglement for mixed states has been studied by some entropic measures such as entanglemnt of formation [1], quantum relative entropy [2] and the degree of entanglement which is defined by entropy minus mutual entropy [3]. In this letter, we will apply the degree of entanglement due to quantum mutual entropy [4], we call it DEM in the sequel, to the entangled state in the Jaynes-Cummings model [5]. Vedral *et al.* defined the degree of entangled state σ as a minimum distance between all disentangled states $\rho \in \mathcal{D}$ such that $E(\sigma) \equiv \min_{\rho\in\mathcal{D}} D(\sigma||\rho)$ where D is any measure of distance between the two states σ and ρ [2]. For an example, one can choose quantum relative entropy as D. Then,

$$E(\sigma) = \min_{\rho\in\mathcal{D}} S(\sigma|\rho), \tag{1}$$

where $S(\sigma|\rho) \equiv tr\sigma(\log\sigma - \log\rho)$ is quantum relative entropy [6,7]. Since this measure has to take a minimum over all disentangled states, it is difficult to calculate analytically for the Jaynes-Cummings model so that we use the DEM defined below. Moreover, there has been no fixed-definition of entanglement measure, though some measures have been defined other than the above measure defined in (1). So we can use the convenient measure which ever we want case-by-case.

Let σ be a state in $\mathfrak{S}_1 \otimes \mathfrak{S}_2$ and ρ_k are the marginal states in $\mathfrak{S}_k$ (*i.e.* $tr_j\sigma = \rho_k\,(k \neq j)$). Then our degree of entanglement due to quantum mutual entropy (DEM) is defined by:

$$I_\sigma(\rho_1,\rho_2) \equiv tr\sigma(\log\sigma - \log\rho_1\otimes\rho_2).$$

Note that the tensor product state $\rho_1\otimes\rho_2$ is one of the disentangled states. This quantity is applied to classify entanglements in [3]. If $\sigma \in \mathfrak{S}_1\otimes\mathfrak{S}_2$ is an entangled pure state, then its von Neumann entropy is equal to 0 (*i.e.*, $S(\sigma) = 0$). Moreover, from the following inequality [8]:

$$|S(\rho_1) - S(\rho_2)| \le S(\sigma) \le S(\rho_1) + S(\rho_2)$$

we have $S(\rho_1) = S(\rho_2)$. Thus we have

$$\begin{aligned} I_\sigma(\rho_1,\rho_2) &= tr\sigma(\log\sigma - \log\rho_1\otimes\rho_2) \\ &= S(\rho_1) + S(\rho_2) - S(\sigma) \\ &= 2S(\rho_1) \end{aligned} \tag{2}$$

Therefore, for entangled pure states, the DEM becomes twice of von Neumann reduced entropy. That is, if we want to know the degree of the entangled pure states, it is sufficient to use von Neumann reduced entropy. However, for entangled mixed states which appear in many cases, we have to use the DEM. From (2), we also find that the entanglement degree in a pure state is bigger than that in a mixed state.

The Jaynes-Cummings model has been studied by many researchers from various points of view [9,10] because this model has a very simple form and can be exactly solvable. One of the most interesting features on this model is the entanglement developed between the atom and the field during the interaction. There have been several approaches to analyze the time evolution in this model, for instance, von Neumann entropy and atomic inversion. In this letter, we will apply the DEM to analyze the entanglement of the Jaynes-Cummings model.

The resonant Jaynes-Cummings model Hamiltonian can be expressed by rotating-wave approximation in the following form

$$\begin{gathered} H = H_A + H_F + H_I, \\ H_A = \frac{1}{2}\hbar\omega_0\sigma_z,\; H_F = \hbar\omega_0 a^* a, \\ H_I = \hbar g(a\otimes\sigma^+ + a^*\otimes\sigma^-), \end{gathered}$$

where g is a coupling constant, $\sigma^{\pm}$ are the pseudo-spin matrices of two-level atom, σ_z is the z-component Pauli spin matrix, a (resp. a^*) is the annihilation (resp. creation) operator of a photon. We now suppose that the initial state of the atom is a superposition of the grounded state and the excited state:

$$\rho = \lambda_0 E_0 + \lambda_1 E_1 \in \mathfrak{S}_A$$

where $E_0 = |1\rangle\langle 1|$, $E_1 = |2\rangle\langle 2|$, $\lambda_0 + \lambda_1 = 1$. Let the field initially be in a coherent state:

$$\omega = |\theta\rangle\langle\theta| \in \mathfrak{S}_F,\ |\theta\rangle = \exp\left(-\frac{1}{2}|\theta|^2\right)\sum_l \frac{\theta^l}{\sqrt{l!}}|l\rangle$$

The continuous map $\mathcal{E}_t^*$ describing the time evolution between the atom and the field for the Jaynes-Cummings model is defined by the unitary operator generated by H such that

$$\begin{aligned} \mathcal{E}_t^* &: \mathfrak{S}_A \longrightarrow \mathfrak{S}_A \otimes \mathfrak{S}_F, \\ \mathcal{E}_t^* \rho &= U_t (\rho \otimes \omega) U_t^*, \end{aligned} \tag{3}$$

$$U_t \equiv \exp\left(-it\frac{H}{\hbar}\right).$$

This unitary operator U_t is written as

$$U_t = \exp(-itH/\hbar) = \sum_{n=0}^{\infty}\sum_{j=0}^{1} E_{n,j}\left|\Phi_j^{(n)}\right\rangle\left\langle\Phi_j^{(n)}\right|, \tag{4}$$

where $E_{n,j} = \exp\left[-it\left\{\omega_0\left(n+\frac{1}{2}\right) + (-1)^j\,\Omega_n\right\}\right]$ are the eigenvalues with $\Omega_n = g\sqrt{n+1}$, called Rabi frequency and $\Phi_j^{(n)}$ are the eigenvectors associated to $E_{n,j}$.

The transition probability which the atom is initially prepared in the excited state and stays at the excited state after the time t is given by

$$\begin{aligned} c(t) &= |\langle n\otimes 2|\, U_t\, |n\otimes 2\rangle|^2 \\ &= \exp\left(-|\theta|^2\right)\sum_n \frac{|\theta|^{2n}}{n!}\cos^2\Omega_n t. \end{aligned}$$



FIG. 1. Transition probability as a function of time t.

FIG.1 shows the transition probability for a coupling constant $g = 1$ and a mean photon number $|\theta|^2 = 5$. This model yields a *dephase* around the time $t_c \approx 1/g$ and shows the damped oscillation with the gaussian envelope. This dampping is caused by the difference and interference of the Rabi frequency Ω_n. This damping phenomenon is often called *Cummings collapses* [10,11]. FIG.1 shows that the atom in this model is the most uncertain state at the time when the transition probability is equal to 0.5, namely $t \approx 3 \sim 7$. So, it can be seen that the system in this model at that time becomes the most correlated. Later the system shows *revival* around $t_r \approx 2\pi|\theta|/g$. The reason of *revival* is considered as a *rephase*, and its revival periodically appears at each $T_k = kt_r$ $(k = 1, 2, 3, \cdots)$. It is also known [10,12] that the system in this model returns most closely to a pure state of the atom around $t_r/2$, during the collapse interval. For details on this model, the readers may refer to the excellent reviews [9,10].

From (3) and (4), the entangled state between the atom and the field follows as

$$\begin{aligned} \mathcal{E}_t^*\rho = {} & e_1(t)\,|2\otimes n\rangle\langle 2\otimes n| \\ & + e_2(t)\,|2\otimes n\rangle\langle 1\otimes n+1| \\ & + e_3(t)\,|1\otimes n+1\rangle\langle 2\otimes n| \\ & + e_4(t)\,|1\otimes n+1\rangle\langle 1\otimes n+1|. \end{aligned} \tag{5}$$

where

$$\begin{aligned} e_1(t) &= \lambda_0 s(t) + \lambda_1 c(t), \\ e_2(t) &= \frac{i}{2}\exp\left(-|\theta|^2\right)(\lambda_1 - \lambda_0)\sum_n \frac{|\theta|^{2n}}{n!}\sin 2\Omega_n t, \\ e_3(t) &= \frac{-i}{2}\exp\left(-|\theta|^2\right)(\lambda_1 - \lambda_0)\sum_n \frac{|\theta|^{2n}}{n!}\sin 2\Omega_n t, \\ e_4(t) &= \lambda_0 c(t) + \lambda_1 s(t), \\ s(t) &= \exp\left(-|\theta|^2\right)\sum_n \frac{|\theta|^{2n}}{n!}\sin^2\Omega_n t. \end{aligned}$$

Taking the partial trace of (5), we have the reduced density operators ρ_t^A and ρ_t^F for the atom and the field, respectively:

$$\begin{aligned} \rho_t^A &= tr_F \mathcal{E}_t^*\rho = e_1(t)\,|2\rangle\langle 2| + e_4(t)\,|1\rangle\langle 1| \\ \rho_t^F &= tr_A \mathcal{E}_t^*\rho = e_1(t)\,|n\rangle\langle n| + e_4(t)\,|n+1\rangle\langle n+1|. \end{aligned}$$

Thus, the degree of the entangled state $\mathcal{E}_t^*\rho$ for Jaynes-Cummings model can be computed as

$$\begin{aligned} I_{\mathcal{E}_t^*\rho}\left(\rho_t^A, \rho_t^F\right) &= tr\mathcal{E}_t^*\rho\left(\log\mathcal{E}_t^*\rho - \log\rho_t^A\otimes\rho_t^F\right) \\ &= S\left(\rho_t^A\right) + S\left(\rho_t^F\right) - S\left(\mathcal{E}_t^*\rho\right) \\ &= -e_1(t)\log e_1(t) - e_4(t)\log e_4(t) \\ &\quad + e_2(t)\log e_2(t) + e_3(t)\log e_3(t). \end{aligned}$$

FIG.2 shows the time development of the DEM for the Jaynes-Cummings model with the mean photon number

$|\theta|^2 = 5$, the coupling constant $g = 1$ and the parameters $(\lambda_0, \lambda_1) = (0.7, 0.3)$.



FIG. 2. The DEM as a function of time t.

From FIG.2, we find the time-development of the DEM for the Jaynes-Cummings model periodically shows the Rabi oscillation, which is an important feature in this model. Moreover, the entanglement degree takes the highest value around the time $t \approx 3 \sim 7$ which corresponds to the collapse interval indicated in FIG.1. Therefore we conclude that the system in this model has the strongest entanglement when the system is the most correlated. This result also coincides with the fact that the entanglement becomes bigger as the entangled state is closer to a pure state, which we also mentioned above in this letter.



FIG. 3. The DEM as functions of parameter λ_0 for T_1(solid curve), T_2(dashed curve), and T_3(dotted curve).

In FIG.3, the entanglement degrees at three revival times T_1, T_2 and T_3 are compared as a function of parameter λ_0. Then we conclude that the entanglement degree takes a higher value as the subscript k in T_k increases, which makes us conjecture the following inequality for every k:

$$I_{\mathcal{E}^*_{T_k}\rho}\left(\rho^A_{T_k}, \rho^F_{T_k}\right) \leq I_{\mathcal{E}^*_{T_{k+1}}\rho}\left(\rho^A_{T_{k+1}}, \rho^F_{T_{k+1}}\right).$$

---

[1] C.H.Bennett, D.P.DiVincenzo, J.A.Smolin and W.K.Wootters, Phys.Rev.A,**54**,3824(1996).

[2] V.Vedral, M.B.Plenio,M.A.Rippin and P.L.Night,Phys. Rev.Lett.,**78**,2275(1997).

[3] V.P.Belavkin and M.Ohya,"Quantum entanglements and entangled mutual entropy",quant-ph/9812082.

[4] M.Ohya, Rep.Math.Phys.**27**,19(1989).

[5] E.T.Jaynes and F.W.Cummings,Proc.IEEE,**51**,89(1963).

[6] H.Umegaki,Kodai Math.Sem.Rep.**14**,59(1962).

[7] M.Ohya and D.Petz, Quantum Entropy and its Use, Sprinver-Verlag, (1993).

[8] H.Araki and E.H.Lieb,Commun.Math.Phys.,18(1970) 160.

[9] H.-I.Yoo and J.H.Eberly,Phys.Rep.**118**,239(1985).

[10] B.W.Shore and P.L.Knight,J.Mod.Optics.**40**,1195(1993)

[11] F.W.Cummings,Phys.Rev.,**140**,A1051(1965).

[12] J.Gea-Banacloche,Phys.Rev.Lett.,**65**,3385(1990)